

# Effectiveness of interventions that support penicillin allergy assessment and de-labelling of patients by non-allergy specialists: a systematic review protocol

Neil Powell<sup>1,2</sup> • Declan Kohl<sup>3</sup> • Shadia Ahmed<sup>4</sup> • Bridie Kent<sup>5</sup> • Jonathan Sandoe<sup>6</sup> • Sarah Tonkin-Crine<sup>7</sup> • Rhys Owens<sup>8</sup> • Jennie Stephens<sup>9</sup> • Mathew Upton<sup>2</sup>

<sup>1</sup>Pharmacy Department, Royal Cornwall Hospital Trust, Truro, Cornwall, UK, <sup>2</sup>School of Biomedical Sciences, University of Plymouth, Plymouth, Devon, UK, <sup>3</sup>School of Biomedical Sciences, University of Leeds and Leeds Teaching Hospitals NHS Trust, Leeds, West Yorkshire, UK, <sup>4</sup>Department of Microbiology, Leeds Teaching Hospitals NHS Trust, Leeds, West Yorkshire, UK, <sup>5</sup>The University of Plymouth Centre for Innovations in Health and Social Care: A JBI Centre of Excellence, Faculty of Health, University of Plymouth, Plymouth, Devon, UK, <sup>6</sup>Leeds Institute of Medical Research, University of Leeds and Leeds Teaching Hospitals NHS Trust, Leeds, West Yorkshire, UK, <sup>7</sup>Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, Oxfordshire, UK, <sup>8</sup>Department of Critical Care, Royal Cornwall Hospital Trust, Truro, Cornwall, UK, and <sup>9</sup>Department of Medicine, Royal Cornwall Hospital Trust, Truro, Cornwall, UK

## ABSTRACT

**Objective:** This review will systematically examine and synthesize the evidence evaluating the effectiveness and safety of interventions that enable non-allergy specialist health care workers to assess allergy risk in patients with reported penicillin allergies and subsequently remove erroneous allergy records.

**Introduction:** The potential benefits of removing erroneous penicillin allergy labels (de-labeling) are wide-ranging. Penicillin allergy assessment and de-labeling is an antibiotic stewardship priority. Delivery of such assessment and de-labeling by non-allergy specialists has been reported in several studies, but the effectiveness and safety have not been formally synthesized. This is a necessary step in the upscaling of penicillin allergy assessment services.

**Inclusion criteria:** This review will consider quantitative studies using appropriate designs. The studies will include adults and pediatric patients who have undergone penicillin allergy assessment and de-labeling by non-allergy specialists in any health care setting.

**Methods:** A range of databases will be searched to identify studies published in English, with no date limit. Unpublished studies and gray literature will also be searched. Title and abstract screening, and assessment of selected full texts against the inclusion criteria will be conducted by at least two independent reviewers. Identified studies will be assessed for methodological quality using standardized critical appraisal instruments. Data will be extracted and categorized using the EPOC taxonomy, and the effectiveness and safety of the intervention will be determined. Where possible, data will be pooled to facilitate meta-analysis. Data from heterogeneous studies will be reported narratively. The GRADE approach for grading the certainty of evidence will be followed.

**Systematic review registration number:** PROSPERO CRD42020219044

**Keywords:** antimicrobial stewardship; penicillin allergy assessment; penicillin allergy de-labeling

*JBI Evid Synth* 2021; 19(0):1–9.

## Introduction

Approximately 6% of the general population in England<sup>1</sup> and 15% of hospital inpatients in England and elsewhere<sup>2–4</sup> have a record of penicillin

allergy. Penicillin-based antibiotics are first-line treatment for many common infections; however, patients with penicillin allergy labels are usually treated with second-line antibiotics.<sup>2</sup> Second-line antibiotics are often more costly,<sup>5–7</sup> less effective in certain clinical circumstances,<sup>1,8–10</sup> and more toxic.<sup>5</sup> Moreover, they are often broader spectrum, which potentially increases a patient's risk of future infections with resistant bacteria.<sup>5,11</sup> Patients with

Correspondence: Neil Powell, neil.powell2@nhs.net

The authors declare no conflict of interest.

DOI: 10.11124/JBIES-21-00075

penicillin allergy records are also associated with exposure to a greater number of antibiotics, increased length of hospital stay,<sup>2,4</sup> and higher hospital readmission rates,<sup>12</sup> all of which increase costs to health care systems.

However, more than 90% of individuals with a penicillin allergy label are not allergic to penicillin.<sup>13</sup> Assessing patients with penicillin allergy labels to identify those who are not allergic to penicillin, and then de-labeling them, has the potential to reduce second-line antibiotic use in favor of penicillin, thereby reducing the unintended consequences associated with second-line antibiotics.

Penicillin allergy assessment of patients with a reported penicillin allergy has traditionally been the domain of allergy experts. Allergy services in the UK and elsewhere are limited,<sup>14</sup> and many hospitals do not have direct access to such services. Furthermore, allergy services do not have capacity to assess and de-label the potentially large number of patients with reported allergies to penicillin. Traditional penicillin allergy testing requires skin testing prior to an oral challenge test, and this is still the main testing method in UK allergy centers. Therefore, penicillin allergy testing is resource-intensive. A less resource-intensive penicillin allergy de-labeling method uses a direct oral penicillin challenge in patients with a history consistent with low risk of future penicillin allergy, forgoing the need for skin testing.

Direct oral challenge testing makes allergy assessment possible outside allergy centers because it is quicker and less resource-intensive than the traditional skin test method. Non-allergy specialist researchers have explored penicillin allergy assessment and de-labeling of hospitalized patients<sup>15,16</sup> and found it to be safe and effective, resulting in increased use of penicillin antibiotics instead of second-line antibiotics with minimal evidence of side effects. Two systematic reviews have confirmed the safety and efficacy of direct oral challenge delivered by allergists and non-allergists as a method of de-labeling adults.<sup>17,18</sup> Other methods delivered by non-allergists, such as skin testing, have also been successfully delivered in the inpatient and outpatient settings.<sup>19,20</sup>

Leading allergists in the US have suggested that every physician needs to obtain an accurate drug intolerance history before avoiding a beta-lactam antibiotic (the broader group name that includes

the penicillin antibiotic group) when it is the drug of choice. These allergists also postulate that addressing unconfirmed beta-lactam allergy on a large scale would lead to a dramatic reduction in the morbidity and mortality associated with unconfirmed beta-lactam allergy and reduce associated health care costs.<sup>21</sup> In September 2020, the American Academy of Allergy, Asthma and Immunology, together with the Infectious Diseases Society of America, wrote to the Centers for Medicare and Medicaid Services to urge US hospitals to include verification of penicillin allergy as part of its mandatory antibiotic stewardship programs.<sup>22</sup> More recently, the World Health Organization recommended antibiotic de-labeling as an effective antimicrobial stewardship strategy.<sup>23</sup> However, knowledge of the epidemiology of penicillin allergy labels and their association with antimicrobial resistance in low- and middle-income countries is sparse, as is the evidence for established antibiotic de-labeling pathways,<sup>24</sup> with the majority of studies occurring in high-income countries.

Penicillin allergy de-labeling is supported by health care workers and accepted by patients. Health care workers in a UK hospital reported frequently encountering patients with penicillin allergy records they believed to be erroneous, and recognized that incorrect penicillin allergy records were a problem.<sup>25</sup> An Australian study demonstrated patient acceptability for oral penicillin challenges to rule out penicillin allergy,<sup>26</sup> while a US study found that patients believed that penicillin allergy testing provided valuable medical information.<sup>27</sup>

Enabling the wider health care workforce to assess patients with penicillin allergy records and to de-label eligible patients is required to ensure penicillin allergy assessment and de-labeling at scale. Lin *et al.*<sup>28</sup> reported on a successful general physician-delivered penicillin allergy de-labeling program in hospitalized patients in the Netherlands. The intervention included physician education, distributing pocket-sized reminder cards, and electronic medical records to prompt physicians to perform the necessary assessment. Maguire *et al.*<sup>29</sup> reported a successful US emergency department physician-delivered, penicillin allergy de-labeling patient pathway. The intervention included the development of a penicillin and cephalosporin test dose procedure guideline, pharmacist-led education, physician-ordered test doses, pharmacist

verification, and nurse administration and post-challenge dose observation.

In this review, we aim to systematically examine the literature to identify and determine the effectiveness of interventions that enable non-allergy specialist health care workers to assess and, where appropriate, de-label adult and pediatric patients with a reported penicillin allergy in any health care setting. The review will also identify and synthesize those components of the interventions that make them safe and effective. A non-allergy specialist is defined as a medical professional whose primary specialization is not in allergy or who has not trained in allergy as part of their specialty.<sup>30</sup>

A preliminary search of PROSPERO, MEDLINE, the Cochrane Database of Systematic Reviews, and *JB I Evidence Synthesis* was conducted and no current systematic reviews on the topic were identified. However, one systematic review by Cooper *et al.*<sup>17</sup> was underway, synthesizing the evidence on the safety and efficacy of de-labeling penicillin allergy in adults using direct oral challenge. This has since been published, and shows that direct oral challenge delivered by both allergists and non-allergists, is safe and effective as a method of de-labeling adults. Our proposed systematic review will be more focused than that of Cooper *et al.*<sup>17</sup> because it will look solely at non-allergists. However, the scope will also be broader because the review will not be limited by health care setting. It will also consider children and adolescents as well as adults, and will include all methods to de-label patients with incorrect penicillin allergy labels. We will narrow our search to non-allergists because we wish to understand the wider frameworks that enable non-allergists to assess penicillin allergy records and safely de-label patients. If this issue is to be tackled at scale, we need to mobilize the non-allergist workforce and teach it how to do this safely.

The findings of this review will inform the development of a complex intervention designed to facilitate and embed penicillin allergy assessment and de-labeling. The intervention will be delivered by non-allergy specialists as part of a secondary care antimicrobial stewardship program in a UK hospital.

The objectives of this systematic review are to i) identify and synthesize the range of interventions and allergy testing methods used by non-allergists to assess reported penicillin allergies and subsequent de-labeling; ii) identify which types of health care

workers have been targeted by interventions to assess and de-label penicillin allergy records (as per the first objective); and iii) determine the effectiveness (increased penicillin antibiotic use) and safety (absence of adverse drug events) of strategies used to deliver non-allergy specialist inpatient de-labeling in hospitalized patients.

## Review question

What is the effectiveness and safety of interventions that facilitate non-allergy specialist health care workers' assessment of adults and pediatric patients with reported penicillin allergy, with subsequent de-labeling of erroneous records?

## Inclusion criteria

### Participants

This review will include studies with any patient (adults, adolescents, or children) who have a penicillin allergy record or self-reported allergy to penicillin, upon direct questioning, in any health care context and from any country.

### Interventions

The review will include studies reporting on penicillin allergy de-labeling using any method (direct de-label, direct oral challenge, skin testing, or oral challenge) by non-allergy specialists, which include, but are not limited to, nurses, pharmacists, and physicians. Penicillin allergy assessment and de-labeling interventions delivered by immunologists or allergy specialists will be excluded.

### Comparators

The comparators will be adults, adolescents, and children who receive usual standard care and do not undergo penicillin allergy assessment. Due to the nature of the intervention, there may not be a comparator group; therefore, studies without a comparator or control group will not be excluded.

### Outcomes

The primary outcomes will be the number of adults, adolescents, or children with a penicillin allergy record who are successfully de-labeled. The secondary outcomes will be i) any measured antimicrobial stewardship impact (eg, antibiotic class prescribed, antibiotic cost, antibiotic side effects, treatment failure, health care associated infections, antibiotic resistant infections); ii) any measured health care

system impact (eg, length of hospital stay, health care resource utilization); and iii) any unintended harm associated with the de-labeling process (eg, anaphylaxis, side effects of antibiotics).

### *Types of studies*

This review will consider both experimental and quasi-experimental study designs, including randomized controlled trials, non-randomized controlled trials, before and after studies, and interrupted time-series studies. In addition, the review will consider analytical observational studies, including prospective and retrospective cohort studies, case-control studies, analytical cross-sectional studies, and descriptive observational study designs. Case reports will be excluded.

### **Methods**

The systematic review will be conducted in accordance with the JBI methodology for systematic reviews of effectiveness,<sup>31</sup> and reported using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist.<sup>32</sup>

The protocol has been registered in PROSPERO (CRD42020219044).

### *Search strategy*

The search strategy will aim to locate both published and unpublished studies. An initial limited search of Embase was undertaken to identify articles on the topic. The text words contained in the titles and abstracts of relevant articles, and the index terms used to describe the articles were used to develop a full search strategy for Embase (Ovid; Appendix I). The search strategy, including all identified keywords and index terms, will be adapted for each included database and/or information source. Backwards and forwards reference searches of all included sources of evidence will be completed to identify additional studies.

Only studies published in English will be included due to a lack of funding for translation services. No date limit will be set for included studies because this is a relatively new antimicrobial stewardship intervention, and studies are only expected to be identified from 2010 onwards.

The databases to be searched from their inception to the present day will include Embase (Ovid), MEDLINE (Ovid), CINAHL (Ovid), PsycINFO, Web of Science, and Cochrane CENTRAL. Sources

of unpublished studies/gray literature will include the World Health Organization Library database, key organization websites and conference proceedings (eg, European Society of Clinical Microbiology and Infectious Diseases, Society for Healthcare Epidemiology of America, Healthcare Infection Society, Infection Prevention Society), registered controlled trial registers, technical or research reports from government agencies, and the British Library (EThOS) Collection of PhD dissertations.

We will contact known experts in the topic regarding any unpublished work and to ensure we have not overlooked relevant literature.

### *Study selection*

Following the search, all identified citations will be collated and uploaded into Endnote Note v.X9.2 (Clarivate Analytics, PA, USA) and duplicates removed. Following a pilot test, titles and abstracts will then be screened by at least two independent reviewers for assessment against the inclusion criteria for the review using RAYYAN software (Qatar Computing Research Institute, Doha, Qatar). Potentially relevant studies will be retrieved in full and their citation details imported into the JBI System for the Unified Management, Assessment and Review of Information (JBI SUMARI; JBI, Adelaide, Australia).<sup>33</sup> The full text of selected citations will be assessed in detail against the inclusion criteria by at least two independent reviewers using RAYYAN software. Reasons for exclusion of papers at full text that do not meet the inclusion criteria will be recorded and reported in the systematic review. Any disagreements that arise between the reviewers at each stage of the selection process will be resolved through discussion or with an additional reviewer. The results of the search and the study inclusion process will be reported in full and presented in a PRISMA flow diagram.<sup>32</sup>

### *Assessment of methodological quality*

Eligible studies will be critically appraised by two independent reviewers at the study level for methodological quality using standardized critical appraisal instruments from JBI for experimental, quasi-experimental, observational, and descriptive studies.<sup>31</sup> Authors of papers will be contacted to request missing or additional data for clarification, where required. Any disagreements that arise will be resolved through discussion or with a third reviewer.

The results of the critical appraisal will be reported in narrative format and in a table.

Studies will not be excluded on the grounds of their risk of bias, but the risk of bias will be reported when presenting the results. The risk of bias judgments will be summarized across different studies for each of the domains listed using the risk of bias graph and the risk of bias summary. Therefore, all studies, regardless of the results of their methodological quality, will undergo data extraction and meta-analysis, where possible.<sup>31</sup>

### Data extraction

Data will be extracted from studies included in the review by two independent reviewers using the standardized data extraction tool from JBI.<sup>31</sup> The extracted data will include specific details about the populations, study methods, interventions, and outcomes of significance to the review objectives and interventions. These data will be categorized using Cochrane's Effective Practice and Organisation of Care taxonomy of health interventions.<sup>34</sup> Authors of papers will be contacted to request missing or additional data, where required.

### Data synthesis

Studies will, where possible, be pooled in statistical meta-analysis using JBI SUMARI.<sup>33</sup> Effect sizes will be expressed as either odds ratios (for dichotomous data) or weighted (or standardized) final post-intervention mean differences (for continuous data), and their 95% confidence intervals will be calculated for analysis. Heterogeneity will be assessed statistically using the standard  $\chi^2$  and  $I^2$  tests, and reasons for heterogeneity will be explored using subgroup and/or sensitivity analyses. Subgroups may include the penicillin allergy de-label method used (eg, direct de-label, direct oral challenge), the setting (inpatient or outpatient), or patient age (adults, adolescents, or children). Statistical analyses will be performed using the random effects model, or if study numbers are small, the fixed effects model.<sup>35</sup> Where statistical pooling is not possible, the findings will be presented in narrative format, including tables and figures, to aid in data presentation. A funnel plot will be generated to assess publication bias if there are 10 or more studies included in a meta-analysis. Statistical tests for funnel plot asymmetry (Egger test, Begg test, Harbord test)<sup>36-38</sup> will be performed where appropriate.

### Assessing certainty in the findings

The Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach for grading the certainty of evidence will be followed,<sup>39</sup> and a Summary of Findings (SoF) will be created using GRADEpro GDT 2020 (McMaster University, ON, Canada). The SoF will present the following information where appropriate: absolute risks for the treatment and control, estimates of relative risk, and a ranking of the quality of the evidence based on the risk of bias, directness, heterogeneity, precision, and risk of publication bias of the review results. The outcomes reported in the SoF will be the proportion of adults, adolescents, or children with a penicillin allergy record who have been successfully de-labeled; any measured antimicrobial stewardship impact; any measured health care system impact; and any unintended harm associated with the de-labeling process.

### Acknowledgments

Katy Oak, outreach librarian, The Knowledge Spa, Royal Cornwall Hospitals Trust, Truro, for her help in developing the search strategy for this review. Chris Johns, information specialist, University of Plymouth, and Natalie King, information specialist, Leeds Institute of Health Sciences, for their technical support.

### Funding

HEE/NIHR ICA Programme Clinical Doctoral Research Fellowship. NIHR300542 September 1, 2020. The funder will have no role in content development.

### References

1. West RM, Smith CJ, Pavitt SH, Butler CC, Howard P, Bates C, *et al.* 'Warning: allergic to penicillin': association between penicillin allergy status in 2.3 million NHS general practice electronic health records, antibiotic prescribing and health outcomes. *J Antimicrob Chemother* 2019; 74(7):2075-82.
2. Powell N, Honeyford K, Sandoe J. Impact of penicillin allergy records on antibiotic costs and length of hospital stay: a single-centre observational retrospective cohort. *J Hosp Infect* 2020;106(1):35-42.
3. Trubiano JA, Smibert O, Douglas A, Devchand M, Lambros B, Holmes, *et al.* The safety and efficacy of an oral penicillin challenge program in cancer patients: a multicenter pilot study. *Open Forum Infect Dis* 2018;5(12):306.



4. Macy E, Contreras R. Health care use and serious infection prevalence associated with penicillin “allergy” in hospitalized patients: a cohort study. *J Allergy Clin Immunol* 2014;133(3):790–6.
5. Blumenthal KG, Shenoy ES, Varughese C, Hurwitz S, Hooper D, Banerji A. Impact of a clinical guideline for prescribing antibiotics to inpatients with reported penicillin or cephalosporin allergies. *J Allergy Clin Immunol* 2015;135(2):AB232-AB.
6. Borch JE, Andersen KE, Bindslev-Jensen C. The prevalence of suspected and challenge-verified penicillin allergy in a university hospital population. *Basic Clin Pharmacol Toxicol* 2006;98(4):357–62.
7. Li M, Krishna MT, Razaq S, Pillay D. A real-time prospective evaluation of clinical pharmaco-economic impact of diagnostic label of ‘penicillin allergy’ in a UK teaching hospital. *J Clin Path* 2014;67(12):1088–92.
8. Blumenthal KG, Shenoy ES, Huang M, Kuhlen JL, Ware WA, Parker RA, *et al.* The impact of reporting a prior penicillin allergy on the treatment of methicillin-sensitive *Staphylococcus aureus* bacteremia. *PLoS One* 2016;11(7):e0159406.
9. Charneski L, Deshpande G, Smith SW. Impact of an antimicrobial allergy label in the medical record on clinical outcomes in hospitalized patients. *Pharmacotherapy* 2011;31(8):742–7.
10. Wood CA, Wisniewski RM. Beta-lactams versus glycopeptides in treatment of subcutaneous abscesses infected with *Staphylococcus aureus*. *Antimicrob Agents Chemother* 1994;38(5):1023–6.
11. Sigona, Steele JM, Miller CD. Impact of a pharmacist-driven beta-lactam allergy interview on inpatient antimicrobial therapy: a pilot project. *J AM Pharm Assoc* 2016;56(6):665–9.
12. Mattingly TJ 2nd, Fulton A, Lumish RA, Williams AMC, Yoon S, Yuen M, *et al.* The cost of self-reported penicillin allergy: a systematic review. *J Allergy Clin Immunol Pract* 2018;6(5):1649–54.e4.
13. National Institute for Health and Care Excellence. Drug allergy: diagnosis and management. Clinical guideline [CG183] [internet]. 2014 [cited 2021 Jan 14]. Available from: <https://www.nice.org.uk/guidance/cg183>.
14. Krishna MT, Huissoon AP, Li M, Richter A, Pillay DG, Sambanthan D, *et al.* Enhancing antibiotic stewardship by tackling “spurious” penicillin allergy. *Clin Exp Allergy* 2017;47(11):1362–73.
15. Chua KYL, Vogrin S, Bury S, Douglas A, Holmes NE, Tan N, *et al.* The penicillin allergy delabeling program: a multicenter whole-of-hospital health services intervention and comparative effectiveness study. *Clin Infect Dis* 2021;73(3):487–96.
16. du Plessis T, Walls G, Jordan A, Holland DJ. Implementation of a pharmacist-led penicillin allergy de-labelling service in a public hospital. *J Antimicrob Chemother* 2019;74(5):1438–46.
17. Cooper L, Harbour J, Sneddon J, Seaton RA. Safety and efficacy of de-labelling penicillin allergy in adults using direct oral challenge: a systematic review. *JAC Antimicrob Resist* 2021;3(1):dlaa123.
18. DesBiens M, Scalia P, Ravikumar S, Glick A, Newton H, Erinne O, *et al.* A closer look at penicillin allergy history: systematic review and meta-analysis of tolerance to drug challenge. *Am J Med* 2020;133(4):452.
19. Englert E, Weeks A. Pharmacist-driven penicillin skin testing service for adults prescribed nonpreferred antibiotics in a community hospital. *Am J Health Syst Pharm* 2019;76(24):2060–9.
20. Wall GC, Peters L, Leaders CB, Wille JA. Pharmacist-managed service providing penicillin allergy skin tests. *Am J Health Syst Pharm* 2004;61(12):1271–5.
21. Vyles D, Macy E. Self-reported beta-lactam intolerance: not a class effect, dangerous to patients, and rarely allergy. *Expert Rev Anti Infect Ther* 2019;17(6):429–35.
22. American Academy of Allergy, Asthma & Immunology. Penicillin allergy testing: AAAAI advocacy priority [internet]. 2020 [cited 2021 Apr]. Available from: <https://www.aaaai.org/about-aaaai/advocacy/penicillin-allergy-testing>.
23. World Health Organization. Antimicrobial stewardship interventions: a practical guide [internet]. 2021 [cited 2021 Jul 2]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/340709/9789289054980-eng.pdf>.
24. Krishna MT, Vedanthan PK, Vedanthan R, El Shabrawy RM, Madhan R, Nguyen HL, *et al.* Is spurious penicillin allergy a major public health concern only in high-income countries? *BMJ Glob Health* 2021;6(5):e005437.
25. Wilcock M, Powell N, Sandoe J. A UK hospital survey to explore healthcare professional views and attitudes to patients incorrectly labelled as penicillin allergic: an antibiotic stewardship patient safety project. *Eur J Hosp Pharm* 2019;26(6):329–33.
26. Trubiano JA, Pai Mangalore R, Baey YW, Le D, Graudins LV, Charles PGP, *et al.* Old but not forgotten: antibiotic allergies in general medicine (the AGM Study). *Med J Aust* 2016;204(7):273.
27. Sundquist, Bowen BJ, Otabor U, Celestin J, Sorum PC. Proactive penicillin allergy testing in primary care patients labeled as allergic: outcomes and barriers. *Postgrad Med* 2017;129(8):915–20.
28. Lin L, Nagtegaal JE, Buijtsels PCAM, Jong E. Antimicrobial stewardship intervention: optimizing antibiotic treatment in hospitalized patients with reported antibiotic allergy. *J Hosp Infect* 2020;104(2):137–43.
29. Maguire M, Hayes BD, Fuh L, Elshaboury R, Gandhi RG, Bor S, *et al.* Beta-lactam antibiotic test doses in the emergency department. *World Allergy Organ J* 2020;13(1):100093.
30. Savic LC, Khan DA, Kopac P, Clarke RC, Cooke PJ, Dewachter P, *et al.* Management of a surgical patient with a label of penicillin allergy: narrative review and consensus recommendations. *Br J Anaesth* 2019;123(1):e82–94.
31. Tufanaru C, Munn Z, Aromataris E, Campbell J, Hopp L. Chapter 3: Systematic reviews of effectiveness. In: Aromataris E, Munn Z, editors. *JBI Manual for Evidence Synthesis* [internet]. Adelaide: JBI; 2020 [cited 2020 Nov 13]. Available from: <https://synthesismanual.jbi.global>.

32. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700-b.
33. Munn Z, Aromataris E, Tufanaru C, Stern C, Porritt K, Farrow J. The development of software and add it to the reference list to support multiple systematic review types: the Joanna Briggs Institute System for the Unified Management, Assessment and Review of Information (JBI SUMARI). *Int J Evid Based Healthc* 2019;17(1):36–43.
34. Cochrane. Effective Practice and Organisation of Care. EPOC taxonomy [internet]. 2015 [cited 2020 Nov 13]. Available from: <https://epoc.cochrane.org/epoc-taxonomy>.
35. Tufanaru C, Munn Z, Stephenson M, Aromataris E. Fixed or random effects meta-analysis? Common methodological issues in systematic reviews of effectiveness. *Int J Evid Based Healthc* 2015;13(3):196–207.
36. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994; 50(4):1088–101.
37. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; 315(7109):629–34.
38. Harbord RM, Egger M, Sterne JA. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Stat Med* 2006;25(20):3443–57.
39. Schünemann H, Brożek J, Guyatt G, Oxman A, editors. GRADE handbook for grading quality of evidence and strength of recommendations using the GRADE approach [internet]. 2013. [cited 2020 Nov 12]. Available from: <https://gdt.gradepro.org/app/handbook/handbook.html>.

## Appendix I: Search strategy

Embase (Ovid) searched October 16, 2020

1. (penicillin adj2 allerg\*).tw.
2. (penicillin adj2 hypersensitiv\*).tw.
3. (penicillin adj2 anaphylaxis).tw.
4. (beta-lactam adj2 allerg\*).tw.
5. (beta-lactam adj2 hypersensitiv\*).tw.
6. (beta-lactam adj2 anaphylaxis).tw.
7. ("betalactam" adj2 anaphylaxis).tw.
8. ("betalactam" adj2 hypersensitiv\*).tw.
9. ("betalactam" adj2 allerg\*).tw.
10. ("\*lactam" adj2 allerg\*).tw.
11. ("\*lactam" adj2 hypersensitiv\*).tw.
12. ("\*lactam" adj2 anaphylaxis).tw.
13. ("antibiotic" adj2 anaphylaxis).tw.
14. ("antibiotic" adj2 hypersensitiv\*).tw.
15. ("antibiotic" adj2 allerg\*).tw.
16. ("antimicrobial" adj2 allerg\*).tw.
17. ("antimicrobial" adj2 hypersensitiv\*).tw.
18. ("antimicrobial" adj2 anaphylaxis).tw.
19. "PENICILLIN DERIVATIVE"/
20. "DRUG HYPERSENSITIVITY"/ or ANAPHYLAXIS/
21. 19 and 20
22. "PENICILLIN ALLERGY"/
23. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 21 or 22
24. "clinical decision tool".tw.
25. "clinical decision making".tw.
26. "clinical assessment tool".tw.
27. direct.tw.
28. challenge.tw.
29. de-label\*.tw.
30. 27 and 28
31. delabel\*.tw.
32. "interview".tw.
33. "antibiotic stewardship".tw.
34. "antimicrobial stewardship".tw.
35. test\*.tw.
36. "allergy assess\*".tw.
37. "oral challeng\*".tw.
38. "ANTIMICROBIAL STEWARDSHIP"/
39. "SKIN TEST"/
40. "PROVOCATION TEST"/



41. “ALLERGY TEST”/
42. “CLINICAL EVALUATION”/
43. ALGORITHM/
44. “RISK ASSESSMENT”/
45. 24 or 25 or 26 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44
46. 23 and 45

No limits set, number of returns 3188.

PROOF